

Laminitis in Dairy Cattle

Like most dairy producers, you've probably seen evidence of laminitis in at least some of your cows at one time or another. Perhaps you've noticed that some of your fresh animals suddenly become lame shortly after calving. Or maybe a few of your older cows have hooves with horizontal grooves and long, curling toes. It's likely that those animals are suffering from laminitis.

Why should you care?

Laminitis is painful for your cows, and each case drains money from your pocket. There is the cost of treating any diagnosed cows and culling cows that otherwise would have remained part of the herd. A cow with laminitis is also reluctant to stand, resulting in lower feed consumption and lower milk yields, with losses of up to 2.8 kg of milk per cow per day. It is also more difficult to detect heats in animals suffering from laminitis because they will be unwilling to stand to be mounted, leading to a lowered reproductive efficiency.

What is laminitis?


Laminitis is an inflammation of the *laminae* and *papillae* in the hoof (see diagram on the right). The layer of tissue inside the hoof, closest to the hoof wall is arranged in regular folds. It is this folded tissue that we are referring to when we talk about the laminae. The equivalent layer of tissue in the sole of the hoof is arranged in irregular folds called papillae. Both the laminae and the papillae act to absorb the shock created by the hoof's impact with the ground. They also produce the horny tissue of the wall and sole. Anytime the blood supply to these tissues is disrupted—through damage to the vessels from internal or external elements—laminitis can result.

There are three phases of laminitis: acute, subclinical and chronic. Acute laminitis is commonly associated with feeding diets too high in concentrates. Sudden changes in diet to higher concentrate rations can cause a condition called acidosis in the rumen. Acidosis causes the release of chemicals into the blood stream, which then travel to the vessels of

the hoof. These chemicals can gradually destroy the vessels supplying the hoof tissues with blood. With restricted blood supply, the laminae and papillae begin to die. The dead tissues swell, increasing the pressure within the rigid hoof wall. The pressure also compresses the living blood vessels—further reducing blood flow—and causing intense pain.

Subclinical laminitis is the most common form of laminitis found in modern dairy operations. This phase of the disease can result from physical injury to the hoof or from damage sustained during acute laminitis. Either way, the layers of tissue between the pedal bone and the hoof wall begin to degenerate, meaning that the pedal bone begins to separate from the hoof wall and sole. At the same time, the damaged tissues are unable to produce horn of the same quality as healthy tissues, leading to a softening of the outer wall and sole of the hoof. Softening of the outer hoof makes it more prone to wear and damage, compounding the problem. Untreated, the disease will progress to chronic laminitis.

During the chronic phase of laminitis, the mechanical damage occurring within the hoof becomes permanent. As the laminae and papillae continue to deteriorate, the pedal bone becomes completely separated from the hoof wall and possibly the sole. Having lost its attachments to the inside of the hoof, the pedal bone rotates so that the front tip points downward, towards the sole. In severe cases, the tip of the pedal bone may actually come through the sole or the entire pedal bone may sink down to the level of the sole.



How can you tell if your cows have laminitis?

The answer to this question depends on the stage of laminitis each cow has developed. A cow that has acute laminitis will likely appear clinically lame, with her feet pulled underneath her body and her back arched. She will often exhibit redness and swelling in the tissues just above the hoof wall and she will be unwilling to stand or walk.

It is only with hoof trimming that signs of subclinical laminitis such as yellowish discoloration of the sole, sole hemorrhages, sole ulcers, double soles and separation of the white line can be seen. This makes regular hoof-trimming important for three reasons:

- trimming allows you to assess how many cows in your herd have developed subclinical laminitis;
- trimming will help cows with subclinical laminitis to recover (as long as the hoof is not continuing to be damaged);
- proper trimming is a good way to help prevent cows from developing subclinical laminitis.

Cows with chronic laminitis may be the easiest to spot. Destruction of the horn-producing tissues disrupts the growth pattern, changing the shape of the hoof. A cow with chronic laminitis will have elongated, flattened, broadened hooves with grooves and ridges on the hoof wall, giving it a rippled appearance. She will also be walking on her heels rather than on her toes and the tips of her toes will curl upwards. This is because the heel continues to grow at a more normal rate while the growth of the front wall is slowed. If her hooves are trimmed, sole ulcers, sole hemorrhages, double soles and yellowish discoloration may be seen.

Risk factors for laminitis

Many producers are aware of the link between concentrate feeding and laminitis, but other important risk factors shouldn't be overlooked. A Swedish study found that first-calf cows were more likely to show signs of sub-clinical laminitis than older cows. In fact, as a cow aged her chances of developing sub-clinical laminitis decreased. Other studies have found that environmental factors can also increase the risk of subclinical laminitis. Steps or curbs greater

Which animals are most at risk for laminitis?

Nutrition related problems:

- ✓ Those fed the highest grain levels:
 - rumen acidosis;
 - large population of gram-negative bacteria releasing endotoxins.
- ✓ Animals close to calving:
 - risk factors related to pregnancy.
- ✓ Fresh cows (2-4 months into lactation):
 - change in diet and environment.

Environment related problems:

- ✓ Animals exposed to uncomfortable environments:
 - concrete floors;
 - high curbs;
 - uncomfortable stalls.

Health related problems:

- ✓ Animals suffering from other diseases
 - especially mastitis.

than 15 cm in height have been linked to laminitis. This may be because cows are forced to transfer their weight onto their hind feet either for a long period of time (if the curb is in front of the feeder) or repeatedly during the day (if the curb is at the entrance to the parlour or the stalls). Inadequate bedding in the stalls can also lead to laminitis problems, possibly due to longer standing times and more half-in, half-out standing. Hard floors, such as concrete, may aggravate problems caused by increased standing times and can predispose cows to developing laminitis. The development of laminitis in cows is not fully understood. Therefore, there is a need for further research into the causes and effective preventative measures for laminitis. Current research at the University of BC is aimed at improving cows' environment to reduce laminitis.

article written by: Erin Bell, University of BC

Cattle Lameness Website

Dr. Paul Greenough, from the Western College of Veterinary Medicine, together with a team of international experts, has recently launched an Internet website dealing with all aspects of cattle lameness. The address is: <http://www.cowdoc.net>